

University of Colorado Hospital - Clinical Laboratory

Stool Collection Guide

GI PCR Panel Plus	Ova and Parasite Exam (O&P)	C. difficile toxin (CDIFTOXPCR)	Fecal Occult Blood
CARY BLAIRING FOR THE PROPERTY OF PROPERTY	TO PERMANDER WITH ADDRESS OF THE STATE OF TH	STERILE TIGHTEN SECURED lane	Approximate Management of the Control of the Contro
Fill to line. Do not overfill!	Fill to line. Do not overfill!		
Fill using a tongue depressor or collection spoon attached to lid.	Fill using a tongue depressor or collection spoon attached to lid.	5-10 mL of sample	Scrape the surface of fecal sample with the sampling
Vigorously shake fluid vials.	Vigorously shake fluid vials.	Formed stools are not accepted.	probe, completely covering the
Pink indicator fluid. Do not use if color fades or turns yellow.	Must send both vials for complete testing.		grooved portion.
Each kit contains: 1 vial with Cary Blair (green lid) 1 empty vial (clear lid) 1 set of collection instructions 1 ziplock bag	1 Zinc-PVA Fixative vial (gray lid)1 Formalin vial (pink lid)	Sterile collection container Specimens in media (such as Cary Blair) are not accepted.	1 FOBT collection vial
Lawson# 68519	Lawson# 13403 & 62149	Lawson# 11053	Lawson# 19753
Stability: 72 hrs ambient or refrigerate	Indefinitely	Deliver to lab immediately	15 days ambient