
2/16/2021 AgileMD | COVID-19 Ambulatory Pathway





Antipyretics

First Choice:

Second Choice:

 Antitussives First line:

Second line:

without cirrhosis

Other medication considerations:

septic shock.

Use COVID patient instructions and work note

Acetaminophen 500 mg 1-2 tablets every 6 hours for patients

Acetaminophen 500 mg q6h PRN for patients with cirrhosis

Ibuprofen 600mg q6h prn for patients without renal disease

Dextromethorphan-guaifenesin 2-20mg/mL oral syrup 10 mL q4h

Codeine-guaifenesin oral liquid 10-100mg/5mL (caution in elderly

Consider not <u>initiating</u> ACE inhibitors and ARBs unless strong

There is no evidence to discontinue these medications

 Theoretrical risk that an increase in ACE2 receptors seen in patients on ACE-I/ARB therapy can lead to worsened infection.

 Avoid systemic steroids per WHO guidelines due to potential harm Can be considered for other reasons such as asthma, COPD or

Give patient COVID video link to supplement patient instructions

If shortness of breath develops, seek immediate medical evaluation

See: COVID-19: Follow-Up Pathway

• Refer to DC video and handouts (English and Spanish)

Benzonatate capsule 100 mg TID prn

patients or those at high risk for delirium)